

**Table A for Final Exam:**

**Table 4. Intercorrelations Among Residualized Change Variables (N = 26)**

Measure	1	2	3	4	5
1. PTSD Checklist	—	.71**	.50**	-.27	-.48*
2. Clinician-Administered PTSD Scale		—	.53**	-.24	-.39
3. Beck Depression Inventory-II			—	-.43*	-.40*
4. COPM-Performance				—	.65**
5. COPM-Satisfaction					—

*Note.* COPM = Canadian Occupational Performance Measure; PTSD = post-traumatic stress disorder.

\* $p < .05$ . \*\* $p < .01$ .

Reference: Speicher, S. M., Walter, K. H., & Chard, K. M. (2014). Interdisciplinary residential treatment of posttraumatic stress disorder and traumatic brain injury: Effects on symptom severity and occupational performance and satisfaction. *The American Journal of Occupational Therapy*, 68(4), 412-421. <https://doi.org/10.5014/ajot.2014.011304>

**Table B for Final Exam:**

Baseline Characteristics of Studied Women				
Characteristics	Controls (30)	Cases (60)	t	p
Age (years)	54.6±3.79	55.5±4.35	-0.964	0.338
BMI (kg/m <sup>2</sup> )	23.01±1.21	29.73±3.33	-13.676	< 0.001
Waist circumference (cm)	76.53±1.87	99.45±11.04	-15.639	< 0.001
Total cholesterol (mmol/L)	6.29±0.89	6.52±0.75	-1.298	0.198
HDL-c (mmol/L)	1.67±0.32	1.43±0.29	3.573	0.001
LDL-c (mmol/L)	4.29±0.60	4.47±0.80	-1.105	0.272
Triglycerides (mmol/L)	1.38±0.47	1.76±0.64	-3.191	0.005
Lipoprotein (a) (g/L)	0.21±0.19	0.23±0.23	-0.589	0.583
HsCRP (mg/L)	0.74±0.27	2.01±1.45	-6.582	< 0.001

*BMI, body mass index; HDL-C, high-density lipoprotein cholesterol; LDL-c, low-density lipoprotein cholesterol; HsCRP, high-sensitivity C-reactive protein.*

Reference: Klisic, A. N., Vasiljevic, N. D., Simic, T. P., Djukic, T. I., Maksimovic, M. Z., & Matic, M. G. (2014). Association between c-reactive protein, anthropometric and lipid parameters among healthy normal weight and overweight postmenopausal women in Montenegro, *Lab Medicine*, 45(1), 12-16. <https://doi.org/10.1309/LMI612RN7AMPEUUL>

## Table C for Final Exam:

Table 2. Completeness of PU documentation in both formats			
PU data elements	E-recording (n=43)	Paper recording (n=39)	Significance (p value)
<b>Descriptions:</b>			
Location	39 (91%)	32 (82%)	0.58
Grade	38 (88%)	28 (72%)	0.02
Size	34 (79%)	19 (49%)	<0.001
Risk assessment	35 (81%)	17 (44%)	0.03
<b>Preventive devices:</b>			
Devices	31 (72%)	21 (54%)	0.04
Repositioning	40 (93%)	34 (87%)	0.13

Note: PU = pressure ulcer

Level of significance: alpha = 0.05

Reference: Tubaishat, A., Tawalbeh, L. I., Azzam, M. A., AlBashtawy, M., & Batiha, A. (2015). Electronic versus paper records: Documentation of pressure ulcer data. *British Journal of Nursing*, 24(6), S30-S37.  
<https://doi.org/10.12968/bjon.2015.24.Sup6.S30>